

Conformational Differences Unfold a Wide Range of Enterotoxigenic Abilities Exhibited by rNSP4 Peptides from Different Rotavirus Strains

Narayan P. Sastri¹, Kiranmayee Pamidimukkala¹, Jagannath R. Marathahalli¹, Suguna Kaza² and C. Durga Rao^{*,1}

¹Department of Microbiology & Cell Biology, ²Molecular Biophysics Unit, Indian Institute of Science, Bangalore 560012, India

Abstract: NSP4 has been recognized as the rotavirus-encoded enterotoxin. However, a few studies failed to support its diarrheagenic activity. As recombinant NSP4 (rNSP4) peptides of different lengths were used in the limited number of studies, a comparison of relative diarrheagenic potential of NSP4 from different strains could not be possible. To better understand the diarrheagenic potential of NSP4 from different strains, in this report we have evaluated the enterotoxigenic activity of the deletion mutant Δ N72 that lacks the N-terminal 72 residues and the biologically relevant Δ N112 peptide which when derived from SA11 rotavirus strain were previously shown to be highly diarrheagenic in newborn mice. Detailed comparative analysis of biochemical and biophysical properties and diarrheagenic activity of the recombinant Δ N72 peptides from seventeen different strains under identical conditions revealed wide differences among themselves in their resistance to trypsin cleavage, thioflavin T (ThT) binding, multimerization and conformation without any correlation with their diarrhea inducing abilities. These results support our previously proposed concept for the requirement of a unique conformation for optimal biological functions conferred by cooperation between the N- and C-terminal regions of the cytoplasmic tail.

Keywords: Rotavirus diarrhea, thioflavin T, diarrheal dose 50 (DD₅₀), nonstructural protein 4, NSP4, virulence, multimerization, viral enterotoxin.

INTRODUCTION

The rotavirus nonstructural protein NSP4, encoded by genome segment 10, is 175 amino acids (aa) in length [1] and has been identified as the viral enterotoxin based on the ability of SA11-NSP4 to induce age-dependent diarrhea in suckling mice [2]. Several studies have revealed that the protein is structurally complex and functionally pleiotropic (Fig. 1A) [1, 3-6]. NSP4 is critical for rotavirus replication, morphogenesis and pathogenesis [7, 8]. It has been reported to exist in multiple forms in the infected cells- as oligomers, higher molecular weight (HMW) complexes [9, 10] and endoplasmic reticulum (ER)- and cytoplasmic membrane-anchored forms [11-14]. Proteolytically-cleaved and secreted forms were also reported [15, 16]. The ER-resident form is anchored through the N-terminal hydrophobic domains and the cytoplasmic tail (CT) of about 131 residues from the C-terminus exhibits all the known important properties associated with the protein including double-layered particle (DLP) binding [11, 12, 17-19] and diarrhea induction [2, 3]. Recent studies have also shown that a pentylsine domain (PD) and amphipathic helical domain (AD) located between residue 55-90 together function as a viroporin domain (VD) [20] (Fig. 1A).

A peptide spanning residue 114-135 was reported to be about 800-fold less efficient in diarrhea induction compared to the full-length protein [2]. However, the sequence from aa 114-135 is highly conserved among different symptomatic

and asymptomatic strains and it alone unlikely determines the optimal diarrhea inducing potential of the protein. Further, a peptide from residue 112-175, secreted from rotavirus infected cells, was reported to induce dose-dependent diarrhea in suckling mice similar to the full-length protein [15]. Recently, we reported that the Δ N72 deletion mutants from simian SA11 and bovine Hg18 strains were about 20-fold more efficient in diarrhea induction in newborn mice than that reported for the full-length protein, and exhibited efficient DLP-binding activity [3].

Rotaviruses exhibit a wide range of DD₅₀ values in newborn mice [21-24]. Rotaviruses can be either symptomatic or asymptomatic, and in some cases NSP4 and a few other viral proteins could cooperatively determine virus virulence [25-32]. Analysis of NSP4 sequences from more than 175 strains failed to identify any residue or motif that could be associated with the virulence phenotype [33]. Mutations at positions 131, 135 and 138 which were reported to result in loss of a porcine rotavirus virulence and diarrhea-inducing ability of the protein [34], did not correlate with the attenuated phenotype of a vaccine strain [32] or the asymptomatic human or feline strains [35-37]. Further, a limited number of studies reported a lack of correlation between virulence of the rotavirus strain and the DD₅₀ of the cognate NSP4 [22, 38, 39], some of which could be due to the use of rNSP4 mutants that varied widely in their length. Based on the analysis of a large number of mutants of the highly diarrheagenic NSP4 from SA11 and Hg18 strains, we recently reported that the biological functions of rNSP4 proteins are dependent on a unique and complex

*Address correspondence to this author at the Department of Microbiology & Cell Biology, Indian Institute of Science, Bangalore 560012, India; Fax: 91-80-23602697; E-mail: cdr@mcbliisc.ernet.in




Fig. (1). (A). Schematic representation of the structural organization of rotavirus NSP4. H1, H2 and H3, N-terminal hydrophobic domains; ECM, Extra Cellular Matrix; ISVD, Interspecies variable domain; DLP, Double-layered particle, (B). Amino acid sequence alignment of the NSP4^{N72} region from different rotavirus strains used in this study. E158K and K158E represent the mutant NSP4 proteins generated by site-directed mutagenesis using PCR from the symptomatic (N136) and asymptomatic (2KD/851) Vellore strains, respectively.

conformation of the cytoplasmic tail [3, 6]. The reported wide variation in diarrhea inducing abilities of a few recombinant proteins could be attributed to improper conformation of the recombinant peptides [22, 38, 39].

This study was undertaken to attempt to resolve the reported inconsistency in the diarrheagenic activity reported for a few rNSP4 peptides, that differed in length, by the analysis of the enterotoxigenic activity of a single polypeptide (Δ N72) of uniform size from a large number of strains. We have evaluated ThT binding, resistance to trypsin, multimerization/oligomerization and conformational properties of the proteins to understand if there is any correlation between any of these properties and their diarrheagenic potential.

MATERIALS AND METHODOLOGY

Viruses and Cells

The rotavirus strains, their G and P serotype/genotype associations and the host from which they were isolated are listed in Table 1. The Vellore neonatal strains were kindly provided by Dr. G. Kang, Christian Medical College, Vellore, India. Except for IS2, 1040, EHP, EC and Vellore strains, viruses were grown in MA104 cells.

Table 1. Rotavirus Strains Used in this Study and their G and P Serotype/Genotype Associations

Strain	Serotype/Genotype	Host	Reference
SA11	G3P[2]	Simian	
RRV	G3P[3]	Rhesus	
NCDV	G6P[1]	Bovine	
Hg18	G15P[21]	Bovine	[40]
EHP	G3P[20]	Murine	
EC	G3P[16]	Murine	
FRV99	G3	Foal	Unpublished
Wa	G1P[8]	Human	
1040	G2P[4]	Human	[41, 42]
IS2	G2P[4]	Human	[41, 42]
ST3	G4P[6]	Human	
I321	G10P[11]	Human	[43]
116E	G9P[11]	Human	[44]
2KD/851	G10P[11]	Human	[45]
99-D/214	G10P[11]	Human	[45]
N136	G10P[11]	Human	[45]
N138	G10P[11]	Human	[45]

The rotavirus strains used in this study and their VP7 serotype and VP4 genotype associations, and the references for the relatively new and less studied viruses are indicated.

References are given only for the relatively less-studied strains.

Cloning, Expression and Purification of Different NSP4 Δ N72 and Δ N112 Peptides

The genomic RNA from EC, EHP, IS2, 1040, I321 and Vellore strains was extracted from the fecal samples and that

from others was isolated from infected cell culture supernatants as described previously [36, 43]. Cloning of the NSP4 gene, its Δ N72 region spanning aa 73 to 175 and generation of the pET22-NH vector for expression of proteins in fusion with an N-terminal His-tag have been described [3]. The Δ N112 region was cloned using strain- and position-specific primers. The nucleotide sequence of Δ N72 and Δ N112 in pBluescript KS+ (pBS) or pET22-NH was determined using T7, M13 forward and/or reverse primers (Macrogen, Korea). All NSP4 Δ N72 and Δ N112 peptides expressed in *E. coli* BL21 (DE3) were highly soluble and were purified by Ni²⁺-NTA-agarose (QIAGEN) chromatography after binding in presence 0.5% NP-40 and washing extensively in its absence [3]. Purity of the proteins was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and mass spectrometry. Molecular masses of the peptides were determined by size exclusion chromatography (SEC) using Sephacryl S-200 column (GE Healthcare) on a Bio-Rad FPLC chromatography system as well as by mass spectrometry using Ultraflex time of flight mass spectrometer (Bruker Daltonics) [3, 46].

Thioflavin T Fluorescence Assay

Δ N72 peptides dialyzed against a buffer containing 10 mM sodium phosphate pH 7.6 and 100 mM NaCl at 100 μ M each of the protein and the dye were used. ThT-binding assays were performed by mixing 50 μ l of 60 μ M protein solution with 450 μ l of 10 μ M ThT. Readings were recorded in a Shimadzu RF-5301 PC spectrofluorometer at 25°C. The excitation wave length was 450 nm, and the emission was monitored between 450 and 600 nm [3, 47].

Determination of Diarrhoeal Dose 50 (DD₅₀) of NSP4 Δ N72 and Δ N112 Peptides

Prior to animal experiments, the ThT binding ability of the Δ N72 peptides was evaluated [3]. Peptides exhibiting high ThT fluorescence were tested between 1 and 100 pmol and those showing highly reduced or lack of ThT binding were evaluated between 50 pmol and 10 nmol in 5-7 day-old BALB/c mouse pups. The recombinant Δ N72 and Δ N112 peptides in 50 μ l of sterile PBS were administered intraperitoneally. DD₅₀ and mean diarrheal scores, on a scale of 1-4, were determined as described [2, 3]. At each dose, 8 mouse pups were used and the experiment was repeated three to four times. The fold efficiency of diarrhea induction of different NSP4 Δ N72 peptides was calculated with reference to the DD₅₀ of SA11 Δ N72 which exhibited the lowest value.

Circular Dichroism (CD) Spectroscopy

Secondary structural differences among different Δ N72 and Δ N112 peptides were examined employing Far UV-CD spectroscopy. The percent α -helical, β -sheet and random conformation contents were determined using the k2d program [48]. CD spectra of the proteins, in 5 mM sodium phosphate buffer pH 7.4 containing 5 mM NaCl, were recorded on a JASCO J-715 spectropolarimeter at a protein concentration of 10 μ M and the molar residue ellipticity was calculated as described [3].

Trypsin Resistance Analysis of Different NSP4 Proteins

Purified Δ N72 peptides were digested with sequencing grade trypsin (Promega) at 37°C, the trypsin-cleaved products were analyzed by Tricine-SDS-PAGE [49] followed by Coomassie Blue staining and the molecular masses of the cleaved products were determined by mass spectrometry as previously described [3]. Relative trypsin resistance on a scale of 0 to 100% was determined by densitometric measurement of intensities of bands corresponding to all the protected fragments of the helical region post 2 hr incubation with respect to control peptide with 75 to 100% being highly resistant and 0 to 25% corresponding to undetectable level of protected fragments.

RESULTS

Thioflavin T Binding Ability of Different NSP4 Δ N72 Peptides

Recently, we have shown that rNSP4 Δ N72 peptides from SA11 and Hg18 were highly diarrheagenic, formed highly ordered higher molecular weight (HMW) complexes, exhibited high α -helical content, thioflavin T binding and resistance to trypsin of the region from residue 73-146 in contrast to a large number of their N- and C-terminal deletion mutants or amino acid (aa) substitution mutants [3, 6]. Mutations in Δ N72 affected the diarrheal-inducing and ThT-binding activities to different extents with DD₅₀ increases ranging between 20-2080-fold (DD₅₀ 0.05 to >10 nmol) [3, 6]. These studies suggested a correlation between optimal ThT binding and efficient diarrheal induction and that efficient ThT binding is dependent on a unique conformation that is significantly affected by aa substitutions throughout the length of the peptide. Further, a specific conformation only in the ordered multimeric forms of SA11- and Hg18-NSP4 Δ N72, but not other HMW complexes of mutant NSP4 Δ N72 peptides, is recognized by ThT. Since NSP4s from different strains exhibit significant aa variations in the flexible C-terminal region of the cytoplasmic tail (CT) (Fig. 1B), it is likely that different NSP4 peptides would differ significantly from each other in their ThT binding property.

The observation that high ThT fluorescence exhibited by SA11- and Hg-18 Δ N72 peptides correlated with their efficient diarrheal-inducing ability suggested that this property can be used to significantly reduce the number of mouse pups required to determine the DD₅₀ of the large number of NSP4 peptides used in this study. Thus a protein showing high ThT binding need not be tested at high concentration and those that exhibit weak binding can be tested only at high concentration.

The Δ N72 peptides from 17 different human (symptomatic and asymptomatic) and animal strains (Table 1) were expressed, purified (Fig. 2A) and their ThT binding property was evaluated. As shown in Fig. (3), only I321-, SA11-, Hg18- and EHP- NSP4 Δ N72 peptides exhibited significant ThT fluorescence and those from all other strains showed either highly reduced or total lack of ThT binding. Of note, whilst NSP4 from the human asymptomatic strain I321 showed about 2.0-3.3-fold more ThT fluorescence than that of Hg18 and SA11, that from other human asymptomatic strains I16E and ST3 exhibited highly reduced

or negligible ThT fluorescence. The peptides from the two murine strains EHP and EC also varied significantly in their ThT binding ability. Thus in contrast to the previous observation with SA11- and Hg18- NSP4 Δ N72 peptides which exhibited high level of ThT binding, the present analysis using similar peptides from a large number of strains revealed that the ThT binding ability varied widely among different peptides.

NSP4 Δ N72 and Δ N112 Peptides Exhibit Similar Pattern in their Diarrheagenic Abilities But with Different DD₅₀ Values

Evaluation of the diarrhea inducing ability of Δ N72 peptides from 17 different strains suggested that different NSP4 Δ N72 peptides used in this study can be identified as either efficient diarrheal inducers represented by SA11, Hg18 and EHP which exhibited a low DD₅₀ of 0.005-0.05 nmol, or inefficient/poor diarrheal inducers represented by other NSP4s exhibiting DD₅₀ > 0.05 nmol (Table 2). Although the efficient diarrheal-inducing activity of the former group correlated, in general, with high ThT fluorescence and spontaneous diarrhea within 30 minutes of administration, the inefficient diarrheal inducers differed widely among themselves without any correspondence between ThT fluorescence and DD₅₀ (Table 2). Further, NSP4s from symptomatic and asymptomatic strains could not be distinguished by their diarrheal inducing abilities. While the NSP4 peptides from human asymptomatic strains ST3, I321, and I16E exhibited DD₅₀ values between 0.07 and 0.5 nmol, those from animal strains and human symptomatic strains (RRV, NCDV, FRV18, EC, and Wa, IS2 and 1040) showed DD₅₀ in the range of 0.75 to >10 nmol. Also, the Δ N72 peptides from the Vellore symptomatic strains (N136 and N138) were about 33-fold less efficient in their ability to induce diarrhea in newborn mouse pups than those from the asymptomatic strains 2KD/851 and 99-D/214. Of note, I321 NSP4 which bound ThT better than SA11 and Hg18 NSP4 Δ N72 peptides, was about 1000-fold less efficient in diarrheal induction (Table 2). Whilst Δ N72 from SA11, Hg18 and EHP induced spontaneous diarrhea within 30-40 min of protein administration at the DD₅₀ value with a mean diarrheal score of 3.2, the inefficient diarrheagenic peptides induced diarrheal stools at or below DD₅₀ values between 1 and 2 hr post administration only after gentle palpation of the abdomen with mean diarrhoeal score of 2.0. Among the latter group, NSP4 from ST3 and NCDV exhibited comparatively lower DD₅₀ (0.07 and 0.075 nmol, respectively). While the peptides from RRV, Wa, equine strain FRV99 and the G2 strain 1040 showed 100-200-fold higher DD₅₀, that from another G2 strain IS2 exhibited 1000-fold higher DD₅₀ than SA11-NSP4 Δ N72 (Table 2). Of significance, the murine EC-NSP4 Δ N72 failed to induce diarrhea even with 10 nmol of the protein (precise DD₅₀ not determined) compared to that of another murine strain EHP (0.05 nmol). Though IS2-NSP4 Δ N72 peptide was very inefficient in diarrheagenic activity (5.0 nmol) among the diarrheagenic Δ N72 peptides, it appears to be significantly better than the full-length proteins from avian strains (41-138 nmol) (Table 2) [23, 39].




Fig. (2). Tricine-SDS-PAGE of NSP4 Δ N72 and Δ N112 polypeptides from different strains. **2A**, Lanes 1-16 represent NSP4 of SA11, RRV, Hg18, NCDV, FRV99, EHP, EC, Wa, S2, ST3, 1040, I321, 116E, N136, 2KD/851 strains and N136-E158K mutant protein. Note that analysis of Δ N72 from only 15 strains and one mutant protein is shown in the figure. Faint bands corresponding to dimmer (D) and tetramer (T) are shown by arrows. **2B**, NSP4 Δ N112 peptides from 6 different strains. M, Pre-stained MW markers (Biorad).

Though Δ N72 from SA11 and Hg18 was very efficient in diarrhea induction, this mutant form of the protein is not detected in the virus infected cells. It is possible that the wide differences observed in the diarrheagenic properties of Δ N72 peptides from different strains could be a manifestation of the unnatural mutation. Since the full-length protein could not be expressed in *E. coli* and that expressed in insect cells was difficult to purify to homogeneity, we expressed and purified the recombinant Δ N112 peptide (Fig. **2B**), which is similar to the biologically active peptide secreted from the virus infected or gene-transfected cells [14], from six different strains (SA11, Hg18, NCDV, I321, EHP and Wa) and determined their DD_{50} in newborn mouse pups. As shown in Table 2, though the pattern of diarrhea inducing ability of the Δ N112 peptides from SA11, Hg18, NCDV, Wa and I321 was very similar to that of the corresponding Δ N72 peptides, the former exhibited high

DD_{50} values than the latter. These results suggest that both Δ N72 and Δ N112 peptides from different strains exhibit very similar pattern in their diarrhea inducing properties, but differ in their relative diarrhea inducing efficiencies. The relatively high DD_{50} values exhibited by the Δ N112 peptides in comparison to those of Δ N72 peptides could be attributed to their lack of the N-terminal amphipathic domain which was shown to potentiate the biological function of the protein [3, 20].

A Single aa Mutation in the Flexible C-Terminus of NSP4 from the Vellore Neonatal G10P[11]-Type Strains Correlates with Virus Virulence without Correspondence with the DD_{50} of the Cognate rNSP4 Δ N72

Recently, G10P [11] strains associated with symptomatic or asymptomatic infections in neonates in Vellore, India, were reported. However, sequence analysis of NSP4 from a

Table 2. Comparative Analysis of Biochemical, Biophysical and Biological Properties of NSP4 Δ N72 Polypeptides from 17 Different Rotavirus Strains

Strain	Preparation	DD ₅₀ (nmol)	Relative ThT Binding*	α -Helix	β -Sheet	Random	% Trypsin Resistance	Percentage Multimer		Percentage Oligomer	
								2.0 mg/ml	0.6 mg/ml	2.0 mg/ml	0.6 mg/ml
SA11	Δ N72	0.005	308.72	61	7	32	75-100	92	90	10	8
	Δ N112	1.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
Hg18	Δ N72	0.006	484.89	62	6	31	75-100	94	91	6	9
	Δ N112	1.5	ND	ND	ND	ND	ND	ND	ND	ND	ND
EHP	Δ N72	0.05	217.42	57	8	35	75-100	77	62	23	38
	Δ N112	>6.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
2KD/851 & 99-D/214	Δ N72	0.06	33.65	41	17	42	75-100	34	29	66	71
E158K mutant	Δ N72	0.06	34.55	40	17	43	75-100	35	30	65	70
ST3	Δ N72	0.07	1.71	59	8	33	50-75	90	85	10	15
NCDV	Δ N72	0.075	2.57	31	16	54	0-25	61	2	39	98
	Δ N112	5.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
1321	Δ N72	0.5	1000	46	22	31	25-50	80	75	20	25
	Δ N112	>6.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
116E	Δ N72	0.5	104.36	57	9	34	25-50	52	40	48	50
RRV	Δ N72	0.75	119.87	48	18	35	50-75	95	72	5	28
1040	Δ N72	0.75	1.47	30	26	44	0-25	0	0	100	100
Wa	Δ N72	1.0	6.40	56	9	35	75-100	38	30	62	70
	Δ N112	>6.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
FRV99	Δ N72	1.0	24.15	40	17	44	ND	ND	ND	98	2
N136 & N138	Δ N72	2.0	87.94	56	9	35	50-75	41	37	59	63
K158E mutant	Δ N72	2.0	ND	ND	ND	ND	ND	ND	ND	ND	ND
IS2	Δ N72	5.0	36.84	36	17	48	25-50	12	5	88	95
EC	Δ N72	>10.0	49.97	39	17	44	75-100	29	21	71	79
Strain	Preparation	DD ₅₀ (nmol)	Reference								
SA11	FL	0.1	[2]								
OSUv	FL	0.175	[33]								
OSUa	FL	>0.5	[33]								
PO13	FL	21	[22], [38]								
	Δ N86	1.0									
	Δ N109	2.0									
TY3	FL	138	[22], [38]								
TY1	FL	108	[22], [38]								
Ch1	FL	41	[22], [38]								

DD₅₀ values of NSP4 Δ N72 and Δ N112 polypeptides from different rotavirus strains from different species in newborn mice are shown. The DD₅₀ values previously reported by others for different NSP4 peptides are also shown. The relative ThT fluorescence emission values are with reference to the lowest-binding Δ N72 from strain 1040. The conformational contents, relative resistance to trypsin digestion, multimerization/oligomerization properties of different NSP4 Δ N72 peptides are provided. Note the asymptomatic strains from which the NSP4 is derived for this study are indicated in yellow shade. The level of efficiency of the proteins in each of the properties is classified into 4 grades in different colour shades. Pink colour refers to either highest efficiency or highest values. Green, gray and red shades denote the decreasing levels of efficiency or the values in the properties. Though efficient diarrhea inducers exhibit high ThT fluorescence, α -helical content, trypsin resistance and efficient multimerization, note a general lack of correlation among the properties of other proteins. FL, full length, ND, Not Determined, OSUv and OSUa refer to OSU virulent and avirulent strains, respectively. E158K and K158E mutant proteins are described in the legend to Fig. (1B).

few isolates failed to differentiate the virulent from the avirulent strains [45]. Sequence analysis, in our laboratory, of NSP4 Δ N72 from two symptomatic Vellore isolates (N136 and N138) for which the sequence was not reported, and two asymptomatic isolates (2KD/851 and 99-D/214) for which the sequence was available, revealed a single aa difference at position 158 (E158K) between the virulent and avirulent pairs. While the symptomatic isolates contained Glu at this position, the asymptomatic strains possessed Lys (Fig. 1B). Only Glu was reported at position 158 in both symptomatic and asymptomatic strains [45]. To understand if the virulence phenotype of the Vellore strains correlates with the diarrheagenic activity of cognate NSP4, the DD₅₀ of Δ N72 peptides from symptomatic and asymptomatic isolates in suckling mice was determined. Unexpectedly, NSP4 Δ N72 peptide from the symptomatic isolates N136 and N138 exhibited DD₅₀ (2.0 nmol) that was about 33-fold higher than that of the asymptomatic isolates 2KD/851 and 99-D/214 (0.06 nmol). Association of the E158K mutation with altered diarrhea induction was further evident from the DD₅₀ values of the E158K and K158E mutant peptides which were indistinguishable from those of the peptides from asymptomatic and symptomatic strains, respectively (Table 2). Of note, N136- Δ N72 consistently showed >50% higher ThT fluorescence of that of 2KD/851, though significantly less than that showed by SA11 and Hg18, (Fig. 3) but was relatively inefficient in diarrhea induction than 2KD/851.

Δ N72 Peptides from Different Strains Differ Significantly in their Resistance to Cleavage by Trypsin

Though ThT is frequently used to detect β -sheet structures in amyloid fibrils and ordered polymeric proteins [50, 51], studies on acetylcholinesterase [52] revealed that ThT binds efficiently to the peripheral ligand binding site which lacks β -sheet structures characteristic of amyloid protein. Though the precise nature of interaction is not understood, these as well as our results [3, 6] suggested that ThT could bind to proteins independent of the amyloid β -sheet structures and that different amino acid substitution and deletion mutant NSP4 Δ N72 peptides differ significantly in their conformation.

The Δ N72 peptide region from different strains contains 17 trypsin cleavage sites that are conserved among the strains (Fig. 1B). Previously, we have shown that single amino acid substitutions in the CT of the highly diarrheagenic SA11-NSP4 Δ N72 affected its diarrhea inducing ability and resistance to trypsin digestion [3]. Since different NSP4s significantly differ in their sequence in the unstructured C-terminal region, it is likely that NSP4s from different strains would vary in their susceptibility to trypsin cleavage. As shown in Fig. (4), the order of trypsin resistance on a scale of 0 to 100 (100 being the highest) was SA11, Hg18, Wa, EHP, EC, 2KD/851 and N136-E158K (75-100%) > ST3, RRV and N136 (50 to 75%) > IS2, I321, 116E (25 to 50%) > NCDV and 1040 (0 to 25%), with the SA11 group being highly resistant and NCDV group being




Fig. (3). ThT fluorescence spectra of different NSP4 Δ N72 proteins. The figure insert shows the ThT fluorescence spectra of NSP4 Δ N72 from the Vellore symptomatic (N136) and asymptomatic (2KD/851) strains and E158K mutant protein generated from N136. Note that the fluorescence spectrum of the mutant protein is identical to that from the asymptomatic strain 2KD/851.




Fig. (4). Relative trypsin Resistance of NSP4 Δ N72 proteins from different strains. The relative resistance to trypsin of the region from residue 72-146 among the different Δ N72 peptides is indicated on a scale of 0-100. M, Top two panels- Prestained MW markers (Biorad), bottom three panels- Low MW range markers (GIBCO-BRL).

extremely susceptible. While Δ N72 from SA11, Hg18 and WA yielded a 9.95 kDa stable fragment, that from the murine strains yielded additional smaller stable fragments (Fig. 4). Further, the non-diarrheagenic EC-NSP4 Δ N72 was as resistant to trypsin as EHP-NSP4 Δ N72. It may be noted that NSP4s from the murine strains exhibit high sequence divergence compared to that of other group A rotaviruses and form a distinct genetic group [22] (Fig. 1B). Of significance, Δ N72 from the symptomatic strain N136 was highly susceptible to trypsin and about 33-fold less diarrheagenic compared to that from the asymptomatic strain 2KD/851 or the N136-E158K mutant in spite of the protein from N136 exhibiting relatively higher ThT fluorescence (Figs. 3, 4 and Table 2).

rNSP4 Δ N72 Peptides from Different Strains Differ in their Conformation and Multimerization Properties

The wide differences in ThT binding and susceptibility to trypsin digestion suggest conformational differences among the recombinant Δ N72 peptides from different strains. CD spectroscopic studies were employed to confirm this prediction. As shown in Fig. (5) and Table 2, all the peptides exhibited highly negative CD spectral values as expected suggesting that they contained the expected coiled coil domain (CCD) and the highly diarrheagenic SA11-, Hg18- and EHP- Δ N72 peptides consistently showed high α -helical content compared to majority of the peptides as observed earlier [3]. Thus while high α -helical content correlated well with the low DD₅₀ values of the efficient diarrhea inducers, the inefficient diarrhea inducers differed significantly among themselves in their conformation contents without

correlation to their relative DD₅₀ values. For example, while NSP4 Δ N72 from Wa, ST3, 116E, N136 and N138 exhibited about 56-58%, that from all other strains showed only 30-48% α -helical content. Generally, proteins exhibiting high α -helicity contained less than 9% β -sheet content and those with low α -helical content showed between 16 and 26%. NCDV-NSP4 Δ N72 differed from all others as it failed to bind ThT and exhibited very high (54%) random conformation content in contrast to that of other proteins which ranged between 32 and 44%. The SA11 Δ N112 peptide contained very low α -helical content compared to the corresponding Δ N72 peptide. Hence the secondary structural contents of Δ N112 peptides from other strains were not evaluated.

Conformational differences appeared to reflect the ordered multimerization ability of the different recombinant proteins. We recently showed that the highly diarrheagenic NSP4 Δ N72 from SA11 and Hg18 existed in HMW complexes in *E. coli*, and that the purified protein formed highly ordered multimers and mutations in the N- or C-terminal regions significantly perturbed the conformation and equilibrium between the multimeric and oligomeric forms to different extents [3, 6]. As shown in Table 1, while the highly diarrheagenic SA11- and Hg18- Δ N72 peptides exhibited efficient multimerization even at low concentration as reported earlier [3], all other Δ N72 peptides differed widely in their multimerization property. Some showed concentration-dependent multimerization (EHP and RRV), whereas others either existed predominantly in oligomeric form or failed to multimerize even at high concentration




Fig. (5). Secondary structure analysis of recombinant NSP4 Δ N72 proteins by Far UV-CD spectroscopy. Purified NSP4s exhibit coiled coil structures as revealed by the highly negative CD spectral values but with significant secondary structural differences among themselves.

(Table 2). Though the rNSP4 Δ N72 from the symptomatic and asymptomatic Vellore strains differing in a single amino acid showed comparable multimerization/oligomerization properties, they differed significantly in their ThT binding, trypsin resistance and conformation contents. NCDV-NSP4 Δ N72 though failed to multimerize at low concentration, about 60% existed as HMW form at high concentration. Interestingly, its diarrhea-inducing ability showed a sharp decline below DD₅₀ value, which probably correlates with the extreme instability of the oligomers at low concentration as seen by the broad peak in SEC (data not shown).

DISCUSSION

Our recent studies based on SA11-and Hg18- Δ N72 peptides and their mutants suggested that efficient diarrhea induction is correlated with high ThT fluorescence, trypsin resistance, α -helical content and/or ordered multimerization [3, 6]. However, the present analysis of different properties of NSP4 Δ N72 peptides on a wider scale revealed that efficient diarrhea inducers exhibited high-level of ThT binding, α -helical content, trypsin resistance, and multimerization. However, no such correlation between the relative DD₅₀ values of the inefficient diarrheagenic peptides and any of the studied properties was observed. (Table 2, Figs. 3-5). Of interest, Δ N72 peptides from asymptomatic strains differed significantly among themselves in their properties. Whilst ST3-NSP4 Δ N72 failed to bind ThT and exhibited DD₅₀ of 0.07 nmol, that of I321, though exhibited high ThT fluorescence, was about 7 fold less efficient in diarrhea induction than the former. 116E Δ N72, though showed DD₅₀ similar to that of I321, exhibited negligible ThT binding. Also, the DD₅₀ of Δ N72 peptides from the asymptomatic strains was either very similar to, or significantly lower than that of some of the symptomatic strains. Further, the NSP4 Δ N72 peptide from the symptomatic Vellore strains N136 and N138 was about 33-

fold less efficient than that of the asymptomatic strains 2KD/851 and 99-D/214 in spite of the former showing higher α -helical content and ThT binding. Linear regression analysis of the data from Table 2 using DD₅₀ as dependent variable and ThT fluorescence, α -helical content, trypsin resistance or percent multimerization as independent variables conclusively revealed a lack of correspondence between diarrheagenic activity and any of the biochemical and biophysical properties (data not shown).

Though it may be argued that Δ N72 peptide does not correspond to the full-length protein or a biologically relevant peptide in the infected cells, it is the longest and highly diarrheagenic peptide from SA11 and Hg18 that could be purified to homogeneity in large quantities from *E. coli* [3]. Present studies using the biologically relevant Δ N112 peptides revealed that both Δ N72 and Δ N112 peptides from different strains exhibit similar pattern of diarrhea inducing properties though differ in the relative range of their DD₅₀ values. The Δ N72 peptide from SA11 and Hg18 is about 20- and 200-fold more efficient in diarrhea induction than that reported for the full-length protein and the secreted Δ N112 peptide [2, 3, 15], respectively in newborn mice and exhibits efficient DLP-binding activity suggesting that this region exhibits optimal biological properties associated with full-length protein. The observation that Δ N112 peptide was less efficient than Δ N72 further supports our previous observation that both the N-terminal AD and C-terminal flexible regions are important for optimal biological functions of the protein [3, 6]. A recent report of the viroporin activity of the N-terminal region from residue 55-90 further supports this observation [20]. Since full-length NSP4 could neither be expressed in *E. coli* due to the presence of hydrophobic domains at the N-terminus nor could that expressed in insect cells be purified to homogeneity [2], different studies used deletion mutants that varied widely in their length and/or the presence or absence of a tag at the N-terminus which might have contributed to

the wide differences in the DD₅₀ values reported in the limited number of studies [2, 3, 15, 23, 39]. Results in our laboratory indicated that presence of a His-tag at the N-terminus of SA11- and Hg18-ΔN72 did not affect its biological function.

Recent studies from our laboratory [3, 6] also suggested that sequence variations observed in the flexible C-terminus compared to the other regions of NSP4s in different strains [33] (Fig. 1B) would affect conformation, ThT binding and biological properties of the recombinant protein. The significant differences in ThT binding, resistance to trypsin digestion, conformation and DD₅₀ values exhibited by the recombinant proteins from the symptomatic and asymptomatic G10P [11] strains that differed at a single amino acid position 158 in the flexible C-terminal region strongly supports this hypothesis. Further, it may be noted that the EC-NSP4 differs from that of EHP at four positions 137, 150, 154 and 161 in the unstructured C-terminus (Fig. 1B). The observation that rNSP4ΔN72 peptides from the murine EHP and EC strains exhibit contrasting diarrhea inducing properties in spite of both viruses being highly virulent in the homologous host strongly suggests a conformational conundrum to the differences in the biological function of the recombinant polypeptides. In the infected cells, NSP4 interaction with other viral [1, 9, 20] and cellular proteins [53-57, 64] might facilitate conformational maturation and the multitude of its biological functions [2, 6, 53, 54, 57-67]. The present studies also suggest that mutations in different NSP4s that could severely affect their biological functions would vary from strain to strain and depend on the overall sequence context of the complete CT, and that rNSP4s, irrespective of their origin, would exhibit DD₅₀ values with inconsistent correlation to the virulence of their virus strains. In this context, our earlier observation that the biological properties of the highly diarrheagenic SA11-NSP4ΔN72 are dependent on a unique and complex conformation in the cytoplasmic tail, mediated by cooperation between the N-terminal amphipathic domain and the extreme C-terminus, is of relevance. The wide variation in conformation among the different peptides, supported by the wide range of differences in susceptibility to trypsin cleavage, ThT binding and multimerization, strongly suggests that unlike the highly enterotoxigenic rNSP4 peptides from SA11 and Hg18, those from majority of the strains fail to attain the proper conformation required for optimal diarrheagenic function.

ACKNOWLEDGEMENTS

Financial support from the Indian Council of Medical Research and the Structural Genomics program under the Genomics Initiative at the Indian Institute of Science funded by the Department of Biotechnology, Indian Council of Medical Research, Government of India is acknowledged. We gratefully acknowledge the use of mass spectrometry and CD facilities in the Molecular Biophysics Unit and the CD facility in the Department of Biochemistry at the Indian Institute of Science. Technical assistance from Mr. Senthil Kumar is acknowledged.

REFERENCES

- [1] Estes MK. In: Knipe DM, Howley PM, Griffin DE, Lamb RA., Martin MA, Roizman B, Straus SE, Eds. *Fields Virology*, 4th ed, vol. 2. Philadelphia; Lippincott Williams and Wilkins 2001; 1747-85.
- [2] Ball JM, Tian P, Zeng CQ-Y, Morris AP, Estes MK. Age-dependent diarrhea induced by a rotavirus nonstructural glycoprotein. *Science* 1996; 272 : 101-4.
- [3] Jagannath MR, Kesavulu MM, Deepa R, *et al.* N- and C-terminal cooperation in rotavirus enterotoxin: novel mechanism of modulation of the properties of a multifunctional protein by a structurally and functionally overlapping conformational domain. *J Virol* 2006; 80: 412-25.
- [4] Taylor JA, O'Brien JA, Yeager M. The cytoplasmic tail of NSP4, the endoplasmic reticulum-localized non-structural glycoprotein of rotavirus, contains distinct virus binding and coiled coil domain. *EMBO J* 1996; 15: 4469-76.
- [5] Deepa R, Rao CD, Suguna K. Structure of the extended diarrhea-inducing domain of rotavirus enterotoxigenic protein NSP4. *Arch Virol* 2007; 152: 847-59.
- [6] Deepa R, Sastri NP, Rao CD *et al.* The flexible C-terminus of the rotavirus nonstructural protein NSP4 is an important determinant of its biological properties. *J Gen Virol* 2008; 89: 1485-96.
- [7] Lopez T, Camacho M, Zayas M, *et al.* Silencing the morphogenesis of rotavirus. *J Virol* 2005; 79: 184-92.
- [8] Silvestri LS, Tortorici MA, Vasquez-Del Carpio R, Patton JT. Rotavirus glycoprotein NSP4 is a modulator of viral transcription in the infected cell. *J Virol* 2005; 79: 15165-74.
- [9] Maass DR, Atkinson PH. Rotavirus proteins VP7, NS28, and VP4 form oligomeric structures. *J Virol* 1990; 64: 2632-41.
- [10] Taylor JA, Meyer JC, Legge MA, *et al.* Transient expression and mutational analysis of the rotavirus intracellular receptor: the C-terminal methionine residue is essential for ligand binding. *J Virol* 1992; 66: 3566-72.
- [11] Bergman CC, Mass D, Poruchynsky M, Atkinson PH, Bellamy AR. Topology of the nonstructural rotavirus receptor glycoprotein NS28 in the rough endoplasmic reticulum. *EMBO J* 1989; 8: 1695-703.
- [12] Chan WK, Au KS, Estes MK. Topology of the simian rotavirus nonstructural glycoproteins (NS28) in the endoplasmic reticulum membrane. *Virology* 1988; 164: 435-42.
- [13] Storey SM, Gibbons TF, Williams CV, Parr RD, Schroeder F, Ball JT. Full-length, glycosylated NSP4 is localized to plasma membrane caveolae by a novel raft isolation technique. *J Virol* 2007; 81: 5472-83.
- [14] Gibbons TF, Storey SM, Williams CV, *et al.* Rotavirus NSP4: Cell type-dependent transport kinetics to the exofacial plasma membrane and release from intact infected cells. *Virol J* 2011; 8: 278.
- [15] Zhang M, Zeng CQ, Morris AP, Estes MK. A functional NSP4 enterotoxin peptide secreted from rotavirus-infected cells. *J Virol* 2000; 74: 11663-70.
- [16] Bugarcic A, Taylor JA. Rotavirus nonstructural glycoprotein NSP4 is secreted from the apical surfaces of polarized epithelial cells. *J Virol* 2006; 80: 12343-9.
- [17] Au K-S, Mattion NM, Estes MK. A subviral particle binding domain on the rotavirus nonstructural glycoprotein NS28. *Virology* 1993; 194: 665- 73.
- [18] O'Brien JA, Taylor JA, Bellamy AR. Probing the structure of rotavirus NSP4: a short sequence at the extreme C terminus mediates binding to the inner capsid particle. *J Virol* 2000; 74: 5388-94.
- [19] Taylor JA, O'Brien JA, Lord VJ, Meyer JC, Bellamy AR. The RER- localized rotavirus intracellular receptor: A truncated purified soluble form is multivalent and binds virus particles. *Virology* 1993; 194: 807-14.
- [20] Hyser JM, Collinson-Pautz MR, Utama B, Estes MK. Rotavirus disrupts calcium homeostasis by NSP4 viroporin activity. *mBio* 2010, 1: e00265-10.
- [21] Burns JW, Krishnaney AA, Vo PT, Rouse RV, Anderson LJ, Greenberg HB. Analyses of homologous rotavirus infection in the mouse model. *Virology* 1995; 207: 143-53.
- [22] Horie Y, Nakagomi O, Koshimura Y, *et al.* Diarrhea induction by rotavirus NSP4 in the homologous mouse model system. *Virology* 1999; 262: 398-407.
- [23] Mori Y, Sugiyama M, Takayama M, Atoji Y, Masegi T, Minamoto N. Avian-to-mammal transmission of an avian rotavirus: Analysis of its pathogenicity in a heterologous mouse model. *Virology* 2001; 288: 63-70.

- [24] Ramig RF. The effects of host age, virus dose, and virus strain on heterologous rotavirus infection of suckling mice. *Microb Pathog* 1988; 4: 189-202.
- [25] Bridger JC, Burke B, Beards GM, Desselberger U. The pathogenicity of two porcine rotaviruses differing in their *in vitro* growth characteristics and gene 4. *J Gen Virol* 1992; 73: 3011-5.
- [26] Broome RL, Vo PT, Ward RL, Clark HF, Greenberg HB. Murine genes encoding outer capsid proteins VP4 and VP7 are not major determinants of host restriction and virulence. *J Virol* 1993; 67: 2448-55.
- [27] Burke B, Desselberger U. Rotavirus pathogenicity. *Virology* 1996; 218: 299-305.
- [28] Hoshino Y, Saif LJ, Kang SY, Sereno MM, Chen WK, Kapikian AZ. Identification of group A rotavirus genes associated with virulence of a porcine rotavirus and host range restriction of a human rotavirus in the gnotobiotic piglet model. *Virology* 1995; 209: 274-80.
- [29] Kirkwood CD, Coulson BS, Bishop RF. G3P2 rotaviruses causing diarrheal disease in neonates differ in VP4, VP7 and NSP4 sequence from G3P2 strains causing asymptomatic neonatal infection. *Arch Virol* 1996; 141: 1661-76.
- [30] Offitt PA, Blavat G, Greenberg HB, Clark HF. Molecular basis for rotavirus virulence: role of gene segment 4. *J Virol* 1986; 57: 46-9.
- [31] Mori Y, Borgan MA, Takayama M, Ito N, Sugiyama M, Minamoto N. Roles of outer capsid proteins as determinants of pathogenicity and host range restriction of avian rotaviruses in a suckling mouse model. *Virology* 2003; 316: 126-34.
- [32] Ward RL, Mason BB, Bernstein DI, et al. Attenuation of a human rotavirus vaccine candidate did not correlate with mutations in the NSP4 gene. *J Virol* 1997; 71: 6267-70.
- [33] Lin SL, Tian P. Detailed computational analysis of a comprehensive set of group A rotavirus NSP4 proteins. *Virus Genes* 2003; 26: 271-82.
- [34] Zhang M, Zeng CQ, Dong Y, et al. Mutations in rotavirus nonstructural glycoprotein NSP4 are associated with altered virus virulence. *J Virol* 1998; 72: 3666-72.
- [35] Chang KO, Kim YJ, Saif LJ. Comparisons of nucleotide and deduced amino acid sequences of NSP4 genes of virulent and attenuated pairs of group A and C rotaviruses. *Virus Genes* 1999; 18: 229-33.
- [36] Jagannath MR, Vethanayagam RR, Reddy BSY, Raman S, Rao CD. Characterization of human symptomatic rotavirus isolates MP409 and MP480 having 'long' RNA electropherotype and subgroup I specificity, highly related to the P6[1], G8 type bovine rotavirus A5, from Mysore, India. *Arch Virol* 2000; 145: 1339-157.
- [37] Oka T, Nakagomi T, Nakagomi O. A lack of consistent amino acid substitutions in NSP4 between rotaviruses derived from diarrheal and asymptotically infected kittens. *Microbiol Immunol* 2001; 45: 173-7.
- [38] Angel J, Tang B, Feng N, Greenberg HB, Bass D. Studies of the role for NSP4 in the pathogenesis of homologous murine rotavirus diarrhea. *J Infect Dis* 1998; 177: 455-8.
- [39] Mori Y, Borgan MA, Ito N, Sugiyama M, Minamoto N. Diarrhea-inducing activity of avian rotavirus glycoproteins, which differ greatly from mammalian rotavirus NSP4 glycoproteins in deduced amino acid sequence in suckling mice. *J Virol* 2002; 76: 5829-34.
- [40] Rao CD, Gowda K, Reddy BSY. Sequence analysis of VP4 and VP7 genes of nontypeable strains identifies a new pair of outer capsid proteins representing novel P and G genotypes in bovine rotaviruses. *Virology* 2000; 276: 104-13.
- [41] Aijaz S, Gowda K, Jagannath HV, et al. Epidemiology of symptomatic rotaviruses in Bangalore and Mysore, India, from 1988 to 1994 as determined by electropherotypes, subgroup and serotype analysis. *Arch Virol* 1996; 141: 715-26.
- [42] Rao CD, Jagannath MR., Varshney BC, Das M, Reddy BSY. In: Kobayashi N. Ed. Genomic diversity and molecular epidemiology of rotaviruses, Trivendrum; Research Signpost 2003; 55-74.
- [43] Das M, Dunn SJ, Woode GN, Greenberg HB, Rao CD. Both surface proteins (VP4 and VP7) of an asymptomatic neonatal rotavirus strain (I321) have high levels of sequence identity with the homologous proteins of a serotype 10 bovine rotavirus. *Virology* 1993; 194: 374-9.
- [44] Gentsch JR, Das BK, Jiang B, Bhan MK, Glass RI. Similarity of the VP4 protein of human rotavirus strain 116E to that of the bovine B223 strain. *Virology* 1993; 194: 424-30.
- [45] Iturriza-Gomara MI, Kang G, Mammen A, et al. Characterization of G10P[11] rotaviruses causing acute gastroenteritis in neonates and infants in Vellore, India. *J Clin Microbiol* 2004; 42: 2541-7.
- [46] Karas M, Hillenkamp F. Laser desorption ionization of proteins with molecular masses exceeding 10000 Daltons. *Anal Chem* 1988; 60: 2299-301.
- [47] Naiki H, Higuchi K, Hosokawa M, Takeda T. Fluorometric determination of amyloid fibrils *in vivo* using the fluorescent dye, thioflavin T. *Anal Biochem* 1989; 177: 244-9.
- [48] Andrade MA, Chacon P, Merelo JJ, Moran F. Evaluation of secondary structure of proteins from UV circular dichroism using an unsupervised learning neural network. *Protein Eng* 1993; 6: 383-90.
- [49] Schagger H, von Jagow G. Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range of 1 to 100 kDa. *Anal Biochem* 1987; 166: 368-79.
- [50] Blanchard BJ, Chen A, Rozeboom LM, Stafford KA, Weigele P, Ingram VM. Efficient reversal of Alzheimer's disease fibril formation and elimination neurotoxicity by a small molecule. *Proc Natl Acad Sci USA* 2004; 101: 14326-32.
- [51] Devlin GL, Chow MKM, Howlett GJ, Bottomley SP. Acid denaturation of α 1-antitrypsin: characterization of a novel mechanism of serpin polymerization. *J Mol Biol* 2002; 324: 859-70.
- [52] Ferrari GV, Mallender WD, Inestrosa NC, Rosenberry TL. Thioflavin T is a fluorescent probe of the acetylcholinesterase peripheral site that reveals conformational interactions between the peripheral and acylation sites. *J Biol Chem* 2001; 276: 23282-7.
- [53] Mirazimi A, Nilsson M, Svensson L. The molecular chaperone calnexin interacts with the NSP4 enterotoxin of rotavirus *in vivo* and *in vitro*. *J Virol* 1998; 72: 8705-9.
- [54] Xu A, Bellamy AR, Taylor JA. Immobilization of the early secretory pathway by a virus glycoprotein that binds to microtubules. *EMBO J* 2000; 19: 6465-74.
- [55] Boshuizen JA, Rossen JW, Sitaram CK, et al. Rotavirus enterotoxin NSP4 binds to the extracellular matrix proteins laminin-beta3 and fibronectin. *J Virol* 2004; 78: 10045-53.
- [56] Parr RD, Storey SM, Mitchell DM, et al. The rotavirus enterotoxin NSP4 directly interacts with the caveolar structural protein caveolin-1. *J Virol* 2006; 80: 2842-54.
- [57] Seo N-S, Zeng CQ-Y, Hyser JM, et al. Integrins α 1 β 1 and α 2 β 1 are receptors for the rotavirus enterotoxin. *Proc Natl Acad Sci USA* 2008; 105: 8811-8.
- [58] Dong Y, Zeng CQ-Y, Ball JM, Estes MK, Morris AP. The rotavirus enterotoxin mobilizes intracellular calcium in human intestinal cells by stimulating phospholipase C mediated inositol 1, 4, 5-triphosphate production. *Proc Natl Acad Sci USA* 1997; 94: 3960-5.
- [59] Tian P, Estes MK, Hu Y, Ball JM, Zeng CQ, Schilling WP. The rotavirus nonstructural glycoprotein NSP4 mobilizes Ca^{2+} from the endoplasmic reticulum. *J Virol* 1994; 69: 5763-72.
- [60] Berkova Z, Crawford SE, Trugnan G, Yoshimori T, Morris AP, Estes MK. Rotavirus NSP4 induces a novel vesicular compartment regulated by calcium and associated with viroplasm. *J Virol* 2006; 80: 6061-71.
- [61] Newton K, Meyer JC, Bellamy AR, Taylor JA. Rotavirus nonstructural glycoprotein NSP4 alters plasma membrane permeability in mammalian cells. *J Virol* 1997; 71: 9458-65.
- [62] Tian P, Ball JM, Zeng CQ, Estes MK. The rotavirus nonstructural glycoprotein NSP4 possesses membrane destabilization activity. *J Virol* 1996; 70: 6973-81.
- [63] Beau I, Cotte-Lafitte J, Geniteau-Legendre M, Estes MK. An NSP4-dependent mechanism by which rotavirus impairs lactase enzymatic activity in brush border of human enterocyte-like Caco-2 cells. *Cell Microbiol* 2007; 9: 2254-66.
- [64] Berkova Z, Crawford SE, Blatt SE, Morris AP, Estes MK. Expression of rotavirus NSP4 alters the actin network organization through the actin remodeling protein cofilin. *J Virol* 2007; 81: 3545-53.
- [65] Martin-Latil S, Cotte-Lafitte J, Beau I, Wuero AM, Geniteau-Legendre M, Servin AL. A cyclic AMP protein kinase A-dependent mechanism by which rotavirus impairs the expression and enzyme activity of brush border-associated sucrase-isomaltase in differentiated intestinal Caco-2 cells. *Cell Microbiol* 2004; 6: 719-31.

- [66] Halaihel N, Lieven V, Ball JM, Estes MK, Alvarado F, Vasseur M. Direct inhibitory effect of rotavirus NSP4(114-135) peptide on the Na(+)-D-glucose symporter of rabbit intestinal brush border membrane. *J Virol* 2000; 74: 9464-70.
- [67] Ousingsawat J, Mirza M, Tian Y, *et al.* Rotavirus toxin NSP4 induces diarrhea by activation of TMEM16A and inhibition of Na⁺ absorption. *Pflugers Arch-Eur J Physiol* 2011; 461: 579-89.

Received: June 20, 2011

Revised: August 18, 2011

Accepted: September 6, 2011

© Sastri *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.